

THE WASHINGTON ENT GROUP

COSMETIC AND ADVANCED FACIAL SURGERY ~ OTOLARYNGOLOGY ~ ENT SURGERY AND SERVICES

Medication List

Section I. The following drugs either contain aspirin and /or have undesirable side effects that may affect your surgery (abnormal bleeding and bruising). These drugs should be avoided for two weeks prior to surgery. This is not every medication known to cause problems. please inform us of any medications that you are currently taking. If you need to take something for relief of minor pain, you may take Tylenol.

| | | |
|------------------------------|----------------------------|--------------------------|
| Advil | Easprin | Pabalate-PF |
| Aleve | Ecotrin | Pamelor |
| Alka-Seltzer | Emprin with codeine | Parnate |
| Alka-Seltzer Plus | Endep | Pepto-Bismol |
| Anacin | Equagesic Tablets | Percodan |
| Anaprox | Etrefon | Persantine |
| An said | Excedrin | Phentermine |
| A.P.C | Feldene | Phenylbutazone |
| Argesic | Fiorinal | Ponstel |
| Arthropan liquid | Fish Oil | Propoxyphene Compound 65 |
| Arthritis Pain Formula | Flagyl | Robaxisal |
| Arthritis Strength Bufferin | Four Way Cold Tablets | Rufen |
| A.S.A | Gelprin | Toradol |
| A.S.A Enseals | Menthcarbamol with Aspirin | Triaminicin |
| Buffets 11 | Micrainin | Trigesic |
| Buffinol | Mobidin | Trilisate |
| Buf-Tabs | Midol | Uracel |
| Butacolidin | Mobigesic | Vanquish |
| Cama Arthritis Pain Reliever | Momentum Backache Formula | Verin |
| Carisoprodol | Motrin | Vitamin E |
| Clinoril | Nalfron | Voltaren |
| Congesprin Chewable | Naprosyn | Zorprin |
| Cope Tablets | Naproxen | |
| Damason P | Nardil | |
| Darvon (all compounds) | Nicobid | |
| Disalcid | Norgesic | |
| Dolprin | Norgesic Forte | |
| Dolobid | Nuprin | |
| | Orudis | |

Section II . The drugs listed below have undesirable side effects that may affect your anesthesia or surgery. Please let us know if your are currently taking any of these medications:

| | | |
|-----------------------|--------------------|---------------------|
| Achromycin | Imipramine HCL | Sinequan |
| Adapin | Isocarboxazid | Surmontil |
| Amitriptyline HCL/MGL | Limnbitrol | Sumycin |
| Aamoxapine | Ludiomil | Tetracycline |
| Anafranil | Maprotiline HCL | Tofranil |
| Asendin | Matulane | Tranlycypromise |
| Aventyl | Medipren | Triavil |
| Capbamazepine | Mysteclin F | Tricyclin |
| Comtrex | Norpramin | Trimipramin maleate |
| Co-Tylenol | Nortriptyline HCL | Vibramycin |
| Desipramin HCL | Novahistine | Vivactil |
| Desyrel | Ornade | Wellbutrin |
| Dilantin | Perphenaxine | Zoloft |
| Doxopin HCL | Phenelzine Sulfate | Zomax |
| Elavil | Procarbazine HOL | Zovirax |
| Etrafor | Protriptyline HCL | |
| Flexeril | Prozac | |

Section III. The following foods contain salicylates that affect blood clotting. They do not need to be omitted completely from your diet, but should be kept to a minimum for two weeks prior to surgery.

| | | |
|-----------|--------------------|-------------------|
| Almonds | Grapes | Plums |
| Apricots | Grape juice | Prunes |
| Berries | Jellies, preserves | Raisins |
| Cherries | Nectarines | Tomatoes |
| Cucumbers | Oranges | Tomatoes products |
| Currants | Peaches | Vinegar |
| Curry | Pickles | Wine |